

12

EUROPEAN PATENT APPLICATION

21 Application number: **87107495.1**

51 Int. Cl.4: **A01N 33/18** , **A01N 25/04**

22 Date of filing: **22.05.87**

30 Priority: **23.05.86 US 867107**

43 Date of publication of application:
16.12.87 Bulletin 87/51

64 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **AMERICAN CYANAMID COMPANY**
1937 West Main Street P.O. Box 60
Stamford Connecticut 06904-0060(US)

72 Inventor: **Morgan, Leonard John**
18 Arundel Drive
Fareham Hampshire(GB)
Inventor: **Bell, Mark**
2 Alan Grove
Fareham Hampshire(GB)

74 Representative: **Wächtershäuser, Günter, Dr.**
Tal 29
D-8000 München 2(DE)

54 **Aqueous suspension concentrate compositions.**

57 The present invention relates to stable aqueous suspension concentrates or aqueous flowable compositions of the low-melting dinitroaniline pesticide, pendimethalin, alone or in combination with secondary pesticide(s) melting at temperatures greater than 70°C or pesticides which are water soluble. Preferred compositions of the invention comprise on a weight to volume basis, about 5.0% to 50.0% pendimethalin; about 0% to 50% of one or more secondary pesticide(s); about 3.0% to 30.0% of coformulants, such as surfactants, dispersing agents, wetting agents, antifreezing agents, antifoaming agents, thickening agents, gums, preservatives and 20.0% to 92.0% water.

The stable compositions of the invention are readily prepared by forming an emulsion of molten pendimethalin in water containing the coformulants, surfactants, dispersing and/or wetting agents, antifoaming agents and suspending agents. This emulsion has an average droplet size in a range of less than 2 microns to about 10 microns, preferably 2 microns to 6 microns. This is then cooled and optionally milled to obtain an average particle size of suspended particles of less than 20 microns, preferably less than 5 microns.

EP 0 249 075 A1

AQUEOUS SUSPENSION CONCENTRATE COMPOSITIONSBACKGROUND OF THE INVENTION

Suspension concentrate pesticidal compositions or aqueous flowable compositions are concentrated suspensions of water-insoluble pesticides and mixtures of pesticides in an aqueous system. The present invention relates to stable such pendimethalin compositions.

These aqueous compositions frequently contain about 10% to 80%, by weight, of a solid pesticide or mixture of solid pesticides, thereby providing a method for handling those pesticides which are relatively water insoluble in an aqueous medium. Since these types of compositions have the desirable characteristics of a thick liquid, they may be poured or pumped. Thus, some of the problems, like dusting that is possible in solid compositions of wettable powders and granulars, are avoided. Further, these aqueous-based concentrates also have the added advantage of not requiring the use of organic solvents, often present in emulsifiable concentrates.

For these reasons, it is desirable to formulate pesticides into suspension concentrates or aqueous flowables. However, such formulations have their own problems such as gelling, caking and settling, as well as problems because of the physical and chemical characteristics of the pesticide or mixture of pesticides. For instance, the dinitroaniline, pendimethalin, is somewhat difficult to formulate and several references have tried to address these formulation problems.

The problems associated with the development of suspension concentrate compositions containing low melting active ingredients, alone or in combination with higher melting active ingredients, are described in German Patent Application DE 3302648 AI. German Patent Application DE 3302648 AI tries to deal with the problems of an aqueous mixed dispersion of a low melting active ingredient in a solvent of phthalic acid C₁-C₁₂ alkyl esters in combination with an aqueous suspension concentrate containing one or more active ingredients as an alternative to a suspension concentrate containing low melting active ingredients, such as pendimethalin[N-(1-ethylpropyl)-2,6-dinitro-3,4-xylidine]. The reason for the alternative approach of that application is the inability to prepare stable suspension concentrates by various techniques, including those of European Patent Application 0 33291 2. That EPO application describes insecticidal suspension concentrate compositions of phosalone and adjuvants which may be prepared with molten insecticide.

Pendimethalin is difficult to formulate not only in a suspension concentrate, but in other forms, as well for several reasons. One is that polymorphic crystals of pendimethalin exist, orange macrocrystals and yellow microcrystals, with the orange form being favored. In formulating pendimethalin in other than suspension concentrates, stabilized pendimethalin had to be used. United States Patents 4,082,537 and 4,150,969 respectively disclose compositions containing either a sodium dialkyl (C₆-C₈) sulfosuccinate or an ethoxylated β -diamine as described. These patents describe the use of sulfosuccinates and β -diamines in order to stabilize pendimethalin's crystal form to the yellow form and to formulate it in a wettable powder composition.

Not only does pendimethalin exist in two crystal forms, but further crystallization occurs when pendimethalin is finally formulated. These formulations often exhibit stability problems related to rapid crystal formation of final product. Very large, elongated crystals (about 3,000 microns in length) are formed in final formulations which result in instability. Thus, formulating compositions wherein these elongated crystals do not develop is crucial to stability of pendimethalin compositions and is necessary to obtain even distribution of active compound for application.

It is an object of the present invention, therefore, to provide stable aqueous suspension concentrate compositions or aqueous flowable compositions of the low-melting dinitroaniline, pendimethalin, either alone or in combination with other pesticides. Although any secondary pesticide may be used, those having higher melting points or pesticides which are water soluble are suited to the compositions of this invention. Further, it is an additional object of the present invention to provide methods for preparing such stable aqueous suspension concentrate compositions or aqueous flowable compositions so that the final compositions do not result in formation of elongated crystals which interfere with processing and active component efficacy for application.

These and other objects will become more apparent by the detailed description of the invention provided herein.

SUMMARY OF THE INVENTION

The present invention relates to stable aqueous suspension concentrate compositions or aqueous flowable compositions comprising pendimethalin, alone or in combination with other pesticides. Although most secondary pesticides can be used, those having melting points greater than 70°C or pesticides which are water soluble are suited in the compositions of the invention. Typically, the compositions of the invention comprise, on a weight to volume basis, about 5.0% to 50.0% pendimethalin; about 0% to 50.0% of one or more secondary pesticide(s); about 3.0% to 30.0% coformulants, as described in more detail hereinbelow; and about 20.0% to 92.0% water.

The stable compositions of the invention are readily prepared by forming an emulsion of molten pendimethalin in water containing the coformulants, surfactants, dispersing and/or wetting agents, antifoaming agents and suspending agents. This emulsion has an average droplet size in a range of less than 2 microns to about 10 microns, preferably 2 microns to 6 microns. This is then cooled and optionally milled to obtain an average particle size of suspended particles of less than 20 microns, preferably less than 5 microns.

DETAILED DESCRIPTION OF THE INVENTION

Preferred compositions of the invention comprise on a weight to volume basis, about 5.0% to 50.0% pendimethalin; about 0% to 50.0% of one or more secondary pesticide(s) having a melting point greater than 70°C or being water soluble; about 3.0% to 30.0% of coformulants, such as surfactants, dispersing agents, wetting agents, antifreezing agents, antifoaming agents, thickening agents, gums, preservatives and 20.0% to 92.0% water.

Coformulants

Pesticides suitable for use in the composition of the present invention include ureas, triazines, imidazolinones, alone or in combination, amongst just a few. Fungicides, insecticides and plant growth regulators which have melting points greater than 70°C and/or possess physical properties which are amenable to the preparation of aqueous suspension concentrate compositions also may be used in the compositions of the present invention.

Additionally, water-soluble pesticides, such as difenzoquat, amine salts, alkali or alkali metal salts of ioxynil, bromoxynil, phenoxy acetic acids, and imidazoliny carboxylic acids such as 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid and the like may readily be incorporated into the stable aqueous suspension concentrate compositions of this invention.

Preferred higher melting (greater than 70°C) components for use in the aqueous suspension compositions of the invention containing pendimethalin include: Isoproturon, [N,N-dimethyl-N'-(4-(1-methylethyl)-phenyl)urea]; Linuron, [N-(3,4-dichlorophenyl)-N'-methoxy-N-methyl urea]; Metoxuron, [N'-(3-chloro-4-methoxyphenyl)N,N-dimethylurea]; Chlortoluron, [N'-(3-chloro-4-methyl-phenyl)-N,N-dimethylurea]; Atrazine, [2-chloro-4-ethyl-amino-6-isopropylamino-1,3,5-triazine]; Imidazolinone herbicides such as 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid and water soluble salts thereof, and the isomeric mixture of methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluate and methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)toluate. Other secondary active components include terbutylazine, 2-tert-butylamino-4-chloro-6-ethylamino-1,3,5-triazine and metolachlor, 2-chloro-6'-ethyl-N-(2-methoxy-1-methylethyl)acet-o-toluidide.

Surfactants (including dispersing agents and/or wetting agents) suitable in the aqueous suspension compositions of the invention containing solid pendimethalin include: ethylene oxide/propylene oxide condensates; alkyl,aryl-and aryl,arylethoxylates and derivatives thereof; lignosulfonates; cresol-and naphthaleneformaldehyde condensates and sulfonates; polycarboxylates and derivatives thereof; and mixtures thereof.

In general, anionic polymerics, such as cresol formaldehyde condensates and their sulfonates, naphthalene formaldehyde condensates and their sulfonates and lignosulfonates have been found to minimize crystal formation during storage and as such, are most preferred.

Suspending agents such as polysaccharide gums like Xanthan gum, guar gum; gum arabic and cellulose derivatives, and the like are suitable for addition to the hot emulsion in amounts of about 0.02% to 3.0%, on a weight to volume basis. These aid in stabilizing the emulsion of a desired droplet size by increasing the viscosity of the emulsion from an initial viscosity of about 100 cps to about 1,000 cps or greater prior to cooling.

Preservatives to prevent microbial spoiling of the compositions of the invention are included as necessary. One example is a 38% solution of formaldehyde. Other preservatives include methyl and propyl parahydroxybenzoate, 2-bromo-2-nitro-propane-1,3-diol, sodium benzoate, glutaraldehyde, O-phenylphenol, benzisothiazolinones, 5-chloro-2-methyl-4-isothiazolin-3-one, pentachlorophenol, 2-4-dichlorobenzylalcohol, mixtures thereof and others known to those in the art. Silicic antifoaming agents are useful in the present compositions.

Antifreezing agents such as ethylene glycol, propylene glycol, other glycols, glycerine or urea may then be added to the resulting aqueous suspension concentrate compositions. Additional surfactants, preservatives and thickening agents, such as clays, precipitated silicas, polyvinyl alcohol, polyvinylpyrrolidone, polyacrylamides and the like, may then be added, as can higher melting active components or a suspension concentrates containing other active components.

Process Of Manufacturing

Stable aqueous suspension concentrate compositions of pendimethalin may be prepared by: emulsifying molten pendimethalin in hot water (50°C to 80°C) containing a surfactant and antifoaming agent to achieve the desired droplet size. Then a suspending agent is added. The resulting hot emulsion is cooled and agitated, allowing the molten material to solidify. The resulting composition may then be milled, if desired, or additional higher melting active components and coformulants, such as antifreezing agents, surfactants, thickeners, preservatives and the like or a preformed suspension concentrate containing one or more active component and conformulants is added. The aqueous suspension concentrate compositions containing pendimethalin in combination with higher melting or water soluble components may then be subjected to additional milling, if desired.

The above method of preparation lends itself to a variety of optional processing steps, such as (1) molten emulsion followed by cooling with no further processing; (2) molten emulsion followed by cooling and optionally adding other active components and coformulants and then milling; (3) molten emulsion in the presence of higher melting active components with concurrent milling followed by cooling; (4) molten emulsion concurrently milled and cooled, then mixing to allow crystallization and standing ("aging") with or without secondary active components and a second milling.

Alternatively, the molten pendimethalin may be dispersed at ambient temperature in a water solution of coformulants, containing if desired, other higher melting active components, followed by milling.

Compositions

Surprisingly, stable aqueous suspension concentrate compositions of pendimethalin, alone or in combination with other active pesticidal components may be prepared by the above methods containing, on a weight to volume basis:

- 5.0% to 50% pendimethalin;
- 0.05% to 1.0% antifoaming agents;
- 2.0% to 20.0% antifreezing agent;
- 2.0% to 20.0% surfactants and mixture of surfactants (wetting and dispersing agents);
- 0.05% to 3.0% thickening agents;
- 0.01% to 1.0% preservatives;
- 0.05% to 2.5% suspending agents;
- and sufficient water to total 100%.

These compositions do not form large, elongated crystals after being processed. Therefore, processing and manufacturing is not halted because of the crystal growths. Further, the compositions are stable without sedimentation of active component in these large (3000 micron) crystals and most importantly, the application of these compositions results in an even dispersibility of active component.

The following examples further illustrate the present invention but are not limitative thereof.

EXAMPLES 1-23Method A

- 5 Preparation of stable aqueous suspension concentrate compositions of pendimethalin, alone or in combination with other higher melting herbicides

10 An aqueous solution containing surfactants and antifoaming agents at temperatures 50°C to 80°C is prepared. Then, the molten pendimethalin (60°C to 80°C) is added and agitated sufficiently to obtain an emulsion having an average droplet size of about 2 microns to 10 microns. This stabilized emulsion is cooled to ambient temperature, allowing the pendimethalin to solidify, whereupon the desired additional coformulants or active components (antifreezing agents, suspending agents, surfactants, pesticides) are added to the resulting aqueous suspension of solid pendimethalin.

15 The resulting aqueous composition is milled to achieve the desired average particle size of suspended particles of less than 20 microns, preferably less than 5 microns; and finally, additional thickening agents, preservatives, or surfactants, as desired, are admixed with the aqueous composition. This is then packaged as the aqueous suspension composition.

Utilizing the above procedure yields the stable aqueous suspension concentrate compositions listed in Table I.

20

25

30

35

40

45

50

55

Example

Composition	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Pendimethalin (unstabilized)	26.0	26.0	23.6	23.0	12.5	26.0	26.0	26.0	26.0	26.0	20.0	20.0	26.0	23.6	40.0
Isoproturon	26.0	26.0	23.6	-	37.5	-	26.0	26.0	26.0	26.0	-	-	26.0	23.6	-
Chlortoluron	-	-	-	30.0	-	-	-	-	-	-	30.0	30.0	-	-	-
Isomeric mixture of methyl 6-(4- isopropyl-4- methyl-5-oxo-2- imidazolin-2-yl)- m-toluate and methyl 2-(4-iso- propyl-4-methyl-5- oxo-2-imidazolin-2- yl)-toluate	-	-	-	-	-	12.5	-	-	-	-	-	-	-	-	-
Na ⁺ cresol- formaldehyde condensate	5.0	-	5.0	-	-	5.0	-	-	-	-	-	-	-	-	-
Na ⁺ cresol- formaldehyde sulphonated condensate	-	3.0	-	3.0	3.0	-	3.0	3.0	3.0	3.0	3.0	3.0	3.0	7.0	5.0
Na ⁺ lauryl sulphate	0.5	0.5	0.5	0.5	0.5	-	0.5	0.5	-	-	-	-	0.5	-	-

TABLE I (Continued)

Composition	Example														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ca ⁺⁺ Lignosulphonate	-	-	-	-	-	2.0	2.0	-	-	-	-	-	-	-	-
Alkyl phenol ethoxylate	-	-	-	-	-	6.0	-	-	-	-	-	-	-	-	-
Urea	8.0	8.0	8.0	8.0	8.0	8.0	-	-	8.0	-	8.0	-	10.0	8.0	-
Precipitated silica	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	-	2.0	2.0	2.0	-
Xanthan gum	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.12	0.12	0.1	0.2
Formaldehyde 38% sol ⁿ	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.5	0.25	0.25	0.25	0.5
Siliconic antifoam	0.1	0.2	0.5	0.25	0.5	0.75	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Ethylene glycol	-	-	-	-	-	-	8.0	8.0	-	8.0	-	8.0	-	-	8.0
Ethylene oxide/pro-pylene oxide condensate	-	-	-	-	-	-	-	-	-	2.0	2.0	2.0	-	-	-
Water	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS	QS

TABLE I (Continued)

Composition	Example								
	16	17	18	19	20	21	22	23	
Pendimethalin	26.0	20.0	20.0	20.0	20.0	20.0	26.0	26.0	
Isoproturon	26.0	-	-	-	-	-	26.0	26.0	
Chlortoluron	-	30.0	30.0	30.0	30.0	30.0	-	-	
Urea	8.0	-	-	-	-	-	-	10.0	
Siliconic antifoam	0.2	0.4	0.5	0.5	0.5	0.5	0.4	0.5	
Xanthan gum	0.2	0.05	-	-	-	0.05	-	0.1	
Formaldehyde 38% solution	0.5	0.125	-	-	-	0.125	-	0.25	
Precipitated silica	-	2.5	2.0	-	-	2.0	2.7	2.0	
Na ⁺ naphthalene-formaldehyde condensate	1.5	-	-	-	-	-	-	-	
Na ⁺ oleoyl methyl tauride	1.5	1.0	-	-	-	-	-	-	
Ethylene oxide-propylene oxide copolymer	-	5.5	2.0	2.0	2.0	3.6	6.0	-	

TABLE I (Continued)

Composition	Example						
	16	17	18	19	20	21	22
Ethylene glycol	-	8.0	8.0	8.0	8.0	8.0	8.0
Na ⁺ cresol-formaldehyde sulphonate condensate	-	-	3.0	3.0	3.0	-	-
Na ⁺ carboxymethyl cellulose	-	-	0.5	-	-	-	-
Polyvinylalcohol	-	-	-	2.0	-	-	-
Polyvinylpyrrolidone	-	-	-	-	2.0	-	-
Na ⁺ Lignosulphonate	-	-	-	-	-	2.0	-
China clay	-	-	-	-	-	-	1.3
Calcium chloride	-	-	-	-	-	-	1.3
Na ⁺ polyacrylate	-	-	-	-	-	-	2.0

Example

Composition	16	17	18	19	20	21	22	23
Propoxylated alkyl-aryl ethoxylate	-	-	-	-	-	-	-	3.0
Water	-	-	-	-	-	-	-	-

-----Sufficient water to total 100%-----

EXAMPLES 24-44Method BPreparation of stable aqueous suspension concentrate compositions of pendimethalin, alone or in combination with other active components

An aqueous solution containing surfactant(s) and antifoaming agents is prepared at a temperature of 50°C to 80°C. Then, the molten pendimethalin (60°C to 80°C) is added to the aqueous solution while agitating sufficiently to obtain an emulsion having an average droplet size of about 2 microns to 10 microns. Sufficient suspending agent is added to this to stabilize the thus-formed emulsion, and this is cooled to ambient temperature, allowing the pendimethalin to solidify, whereupon the additional coformulants, as desired, are admixed to the emulsion. This can then be packaged.

Further, a suspension of a pesticide having a melting point greater than 70°C is prepared and milled to a suitable average particle size (i.e. less than 20 microns, preferably less than 5 microns) or an aqueous solution containing the desired amount of a water-soluble pesticide is prepared.

Either one of these is then admixed with the suspension concentrate composition of pendimethalin prepared hereinabove. Finally, additional thickening agents, preservatives or surfactants, as desired, are added, and this is then packaged as the mixed aqueous suspension concentrate composition.

Utilizing the above procedure yields the stable aqueous suspension concentrate compositions listed in Table II.

TABLE II

Composition	Example			
	24	25	26	27
Pendimethalin (unstabilized)	23.6	23.6	23.6	23.6
Isoproturon	23.6	23.6	23.6	23.6
Na ⁺ cresol-formaldehyde sulphonated condensate	-	3.0	4.1	3.0
Na ⁺ cresol-formaldehyde condensate	5.0	-	-	-
Alkylphenol ethoxylate	-	-	-	-
Na ⁺ oleoyl methyl tauride	-	0.65	-	-
Na ⁺ lauryl sulphate	0.5	-	-	0.5
Urea	8.0	8.0	8.0	-
Ethylene glycol	-	-	-	8.0
Blend of polyalkylene glycol ether and polyoxy-ethylene alkylaryl ether	-	-	0.7	-
Siliconic antifoam	0.25	0.5	0.5	0.5
Silica	2.0	2.0	-	2.0
Xanthan gum	0.1	0.1	0.2	0.1
Formaldehyde 38% soln	0.25	0.25	0.5	0.25
Water	QS	QS	QS	QS

TABLE II (Continued)

Composition	Example				
	29	30	31	32	33
Pendimethalin (unstabilized)	20.0	20.0	20.0	26.0	30.0
Chlortoluron	30.0	30.0	-	-	-
Isomeric mixture of methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazol in-2-yl)-m-toluate and methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazol in-2-yl)-toluate	-	-	12.5	12.5	-
Ammonium 2-(4-isopropyl-4-methyl-5-oxo-2-imidazol in-2-yl)-3-quinolinecarboxylic acid (solution)	-	-	-	-	5.0
Na ⁺ cresol-formaldehyde sulphonated condensate	3.0	3.6	-	-	-
Na ⁺ cresol-formaldehyde condensate	-	-	5.0	3.0	-
Triethanolamine salt of polyarylaerylethoxylate phosphate	-	-	-	1.3	-
Na ⁺ Lignosulphonate	-	-	-	-	4.0
Ca ⁺⁺ Lignosulphonate	-	-	2.0	-	-
Alkylphenol ethoxylate	-	-	8.0	6.0	-
Na ⁺ lauryl sulphate	0.5	-	-	-	-
Urea	8.0	8.0	8.0	-	-

TABLE II (Continued)

Composition	Example				
	29	30	31	32	33
Ethylene glycol	-	-	-	5.0	8.0
Siliconic antifoam	0.5	0.1	0.5	0.4	0.5
Silica	2.0	0.1	2.0	2.0	0.75
Xanthan gum	0.1	0.2	0.1	0.1	0.1
Formaldehyde 38% soln	0.25	0.5	0.25	0.25	0.25
Glacial acetic acid (to pH 7.4)	-	-	-	-	✓
Blend of polyalkylene glycol ether and polyoxy- ethylene alkylaryl ether	-	0.55	-	-	-
Water	QS	QS	QS	QS	QS

TABLE II (Continued)

Composition	Example							
	34	35	36	37	38	39		
Pendimethalin (unstabilized)	33.0	40.0	40.0	40.0	40.0	33.0		
Na ⁺ cresol-formaldehyde sulphonated condensate	-	4.2	-	-	-	-		
Na ⁺ cresol-formaldehyde condensate	-	-	4.7	-	-	4.2		
Triethanolamine salt of polyarylethoxylate phosphate	3.4	-	-	-	-	-		
Na ⁺ Lignosulphonate	-	-	-	4.8	-	-		
Urea	-	-	-	-	-	13.3		
Ethylene glycol	8.0	8.0	-	-	5.0	-		
Siliconic antifoam	0.1	0.5	0.9	0.3	1.0	0.3		
Xanthan gum	0.1	0.1	0.1	0.1	0.2	0.16		
Formaldehyde 38% soln	0.25	0.25	0.25	0.25	0.5	0.4		
Polycarboxylate derivative	-	-	-	-	3.0	-		
Water	QS	QS	QS	QS	QS	QS		

TABLE II (Continued)

Composition	Example			
	40	41	42	43
Pendimethalin (unstabilized)	27.3	30.0	27.3	30.0
Atrazine	18.2	20.0	18.2	20.0
Na ⁺ cresol-formaldehyde sulphonated condensate	3.4	1.65	3.4	1.65
Triethanolamine salt of polyarylethoxylate phosphate	-	-	-	1.9
Urea	5.8	-	-	5.0
Ethylene glycol	-	5.6	5.2	5.0
Blend of polyalkylene glycol ether and polyoxyethylene alkylaryl ether	1.34	1.26	1.2	-
Siliconic antifoam	0.9	0.35	0.9	0.16
Silica	-	0.5	-	0.4
Xanthan gum	0.2	0.1	0.2	0.1
Formaldehyde 38% soln	0.5	0.25	0.5	0.25
Water	QS	QS	QS	QS

EXAMPLES 45 and 46

5

Method CPreparation of stable aqueous suspension concentrate compositions

- 10 An aqueous dispersion of surfactants, antifoaming and antifreezing agents, containing, if desired, a solid active component having a melting point greater than 70°C, is prepared or a water soluble active component is prepared at ambient temperatures. The molten pendimethalin (60°C to 80°C), with or without additional surfactants, is then added to the agitated aqueous mixture. This resulting aqueous mixture is milled to achieve the desired average particle size of suspended solids, less than 20 microns, preferably
- 15 less than 5 microns, and additional thickening agents, suspending agents, preservatives, antifreezing agent and surfactants, as desired, are admixed to the milled composition. This is then packaged as the resulting stable aqueous suspension concentrate composition.

Utilizing the above procedure yield the stable aqueous concentrate compositions listed in Table III.

20

25

30

35

40

45

50

55

TABLE III

Composition	Example	
	45	46
Pendimethalin (unstabilized)	26.0	26.0
Isoproturon	26.0	26.0
Na ⁺ cresol-formaldehyde sulphonated condensate	3.0	-
Polyarylpolyoxyethylene phosphate, acid form	-	5.0
Siliconic antifoam	0.2	0.2
Xanthan gum	0.2	0.12
Formaldehyde 38% solution	0.5	0.5
Ethenediol (Ethylene glycol)	8.0	-
Water	QS	QS

EXAMPLES 47 and 48Methods A and E

Preparation of stable aqueous suspension concentrates of pendimethalin via the method of the invention versus the method of milling while cooling an emulsion (Method E)

An aqueous suspension concentrate composition of pendimethalin was prepared according to Examples 1-23, Method A. As a comparison, a suspension concentrate of pendimethalin was prepared according to the description of EPO Application 033291.2.

A mixture of hot water (575 cc), ethylene glycol (50 g) and an anionic surface-active agent (a mixture of the monophosphate and the diphosphate of tristerylphenol with a polyoxyethylene of 18 oxyethylene units, neutralized with triethanolamine, marketed under the tradename of Soprophor FL® by Rhone-Poulenc) (50 g) is vigorously agitated while 400 g of pendimethalin are added. This is then ground in a dyno mill with a jacket for rapid cooling, resulting in a mixture leaving the mill at 24°C and having a particle size of 98% less than 5 microns, indicating the formation of a suspension concentrate.

A Xanthan gum biopolymer of a heteropolysaccharide type (1.5 g), produced by fermentation of Xanthomonas campestris on carbohydrates (tradename Rhodopol XB 23 marketed by Rhone-Poulenc), is added.

Table IV summarizes the stability observations of the two compositions. The milling method does not avoid the appearance of elongated crystals even upon two short periods of storage, one three day storage at 15°C and one three day storage at 28°C.

TABLE IV

Method A - Method of the Invention			Method E - Milling while cooling*		
Composition	% w/v	Result	Composition	%w/w	Result
Pendimethalin	40	(1) After six weeks at 280C, 95% of the particles have an average particle size of less than 15 x 4 microns.	Pendimethalin	40	Two tests: (1) Large crystals particle size up to 80 x 5 microns appeared after three days at 150C and also when stored at three days at 280C.
Suspending agent Xanthan gum (added prior to cooling)	0.02		Suspending agent Xanthan gum (added after milling and cooling)	0.15	
Surfactants			Surfactants		
Sulfonated cresol formaldehyde condensate	4.0		mixture of mono and di phosphate of tristyrylphenol neutralized with triethanolamine (Soprophor®FL)	5.0	
Polyethoxylated-polymethylmethacrylate	3.0		Antifreeze		
Urea	5.0		Ethylene glycol	5.0	
Antifoam					
Siliconic	0.5				
Preservative					
Benzisothiazolone	0.1				
*Trademark of Soprosioie, Division of Rhone-Poulenc.			*Particle size found in initial suspension concentrate composition is 98% less than five (5) microns.		

EXAMPLES 49-50Compositions containing active compound of the invention

An aqueous suspension concentrate composition of pendimethalin with a secondary pesticide was prepared according to Examples 1-23 containing the following components:

10

		(% w/v)	
		Formula X	Formula Y
15	Pendimethalin	23.6	20.0
	Isoproturon	23.6	-
	Chlortoluron	-	-
20	Na cresol-formaldehyde Sulphonated condensate surfactant	4.1	4.14
	Polycarboxylate surfactant	-	1.64
25	Blend of polyalkylene glycol ether and polyoxyethylene alkyl aryl ether	0.71	-
30	Urea	8.0	8.0
	Xanthan gum	0.2	0.2
	30% siliconic antifoam	0.5	0.3
35	Benzisothiazolinone	-	0.033
	Methyl paraben	0.1	-
	Propyl paraben	0.05	-
40	Water	To 100%	100%

EXAMPLES 51-52Compositions using hot emulsion double milling process

45

A hot emulsion of molten pendimethalin is prepared as in Examples 1-23 wherein hot pendimethalin is added to the hot solution of surfactant whilst mixing at high shear. The temperature is 50°C to 80°C with a particle size of 2 μ to 5 μ . The hot emulsion is then milled through a Dyno-Mill and exits as a shattered crystalline form (temperature into mill is about 65°C, and temperature exiting is about 20°C to 25°C). This mixture is allowed to "age" to allow orange polymorph crystal conversion. Following conversion, usually 0.5 to 48 hours, a second milling in a Dyno-Mill occurs. (In the event another active pesticide is added, it is added into the aging period stage). Once milled a second time, final coformulants are added.

The following compositions are prepared according to the above procedures.

55

		<u>Concentrate</u>
		<u>% w/w</u>
5	Pendimethalin technical	40.0
	Soprophor® FL surfactant	5.0
10	Silnaolapse® 500	0.5
	Propylene glycol	7.0
	Water	To 100%

15 This is formulated into the following:

		<u>8/1</u>
	Concentrate from above	1000
20	Rodopol 2% gel	70
	Water	To 100%

		<u>% w/v</u>
25	Concentrate from above	75.0
	Atrazine technical	20.0
	Soprophor®FL	1.25
30	Propylene glycol	1.25
	Water	To 100%

35 Claims

1. An aqueous suspension concentrate composition comprising, on a weight to volume basis: about 5.0% to 50.0% pendimethalin; about 0% to 50.0% of one or more secondary pesticide(s); about 3.0% to 30.0% coformulants; and about 20.0% to 92.0% water.

40 2. A composition according to Claim 1, wherein said secondary pesticide(s) is water soluble or has a melting point greater than 70°C.

3. A composition according to Claim 2, wherein said coformulants are surfactants; dispersing agents; wetting agents; suspending agents; antifreezing agents; antifoaming agents; thickening agents; and preservatives.

45 4. A composition according to Claim 3, wherein said surfactants, dispersing agents and wetting agents are ethylene oxide/propylene oxide condensates; alkyl,aryl-and aryl,aryl-ethoxylates and derivatives thereof; lignosulfonates; cresol-formaldehyde condensates and sulfonates; naphthalene-formaldehyde condensates and sulfonates; polycarboxylates and their derivatives; and mixtures thereof.

50 5. A compositions according to Claim 4, wherein said suspending agents are polysaccharide gums and cellulose derivatives.

6. A composition according to Claim 5, wherein said polysaccharide gums are Xanthan gum, guar gum, gum arabic, and mixtures thereof.

7. A composition according to Claim 6, wherein said antifreezing agents are ethylene glycol, propylene glycol, glycerine, urea, and mixtures thereof.

55 8. A composition according to Claim 7, wherein said thickening agents are clays, precipitated silicas, polyvinyl alcohol, polyvinylpyrrolidone, polyacrylamides, and mixtures thereof.

9. A composition according to Claim 8, comprising, on a weight to volume basis: 5.0% to 50.0% pendimethalin; 0% to 50.0% of a pesticide with a melting point greater than 70°C or is water soluble; 2.0% to 20.0% surfactants; 0.05% to 2.5% suspending agents; 2.0% to 15.0% antifreezing agents; 0.05% to 2.0% thickening agent; and 0.05% to 2.5% preservatives.

5 10. A composition according to Claim 9, wherein said antifoaming agent is a silicic antifoaming agent.

11. A composition according to Claim 10, wherein said preservative is a 38% formaldehyde solution, methyl or propyl parahydroxybenzoate, 2-bromo-2-nitro-propane-1,3-diol, sodium benzoate, glutaraldehyde, O-phenylphenol, benzisothiazolinones, 5-chloro-2-methyl-4-isothiazolin-3-one, pentachlorophenol, 2,4-dichlorobenzylalcohol, or mixtures thereof.

10 12. A composition according to Claim 10, wherein said pesticide with a melting point greater than 70°C is Isoproturon, [N,N-dimethyl-N'-(4-(1-methylethyl)phenyl)urea]; Linuron, [N3,4-dichlorophenyl]-N'-methoxy-N'-methyl urea; Metoxuron, [N'-(3-chloro-4-methoxyphenyl)-N,N-dimethylurea]; Chlortoluron, [N'-(3-chloro-4-methylphenyl)-N,N-dimethylurea]; Atrazine, [2-chloro-4-ethylamino-6-isopropyl amino-1,3,5-triazine], ter-butylazine, 2-tert-butylamino-4-chloro-6-ethylamino-1,3,5-triazine and metolachlor, 2-chloro-6'-ethyl-N-(2-methoxy-1-methylethyl)ace-o -toluidide; Imidazolinone herbicides such as 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid and water soluble salts thereof, or the isomeric mixtures of methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluate and methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-toluate.

13. A composition according to Claim 10, comprising, on a weight to volume basis: 30.0% to 40.0% pendimethalin; 3.0% to 5.0% sodium cresol-formaldehyde condensate or sodium cresol-formaldehyde sulphonated condensate; 5.0% to 10.0% ethylene glycol or urea; 0.1% to 1.0% antifoaming agent; 0.1% to 0.3% Xanthan gum; 0.2% to 1.0% preservative; and 48.3% to 68.3% water.

14. A composition according to Claim 11, comprising, on a weight to volume basis: 40.0% pendimethalin; 5.0% sodium cresol-formaldehyde sulphonated condensate; 8.0% ethylene glycol; 0.5% 25 silicic antifoaming agent; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution and 51.4% water.

15. A composition according to Claim 12, comprising, on a weight to volume basis: 27.3% pendimethalin; 18.2% atrazine; 3.4% sodium cresol-formaldehyde sulphonated condensate; 5.8% urea; 1.34% polyalkylene glycol ether and polyoxyethylene alkylaryl ether; 0.9% silicic antifoaming agent; 0.2% xanthan gum; 0.5% of a 38% formaldehyde solution; and 48.9% water.

30 16. A composition according to Claim 12, comprising, on a weight to volume basis: 23.6% pendimethalin; 23.6% isoproturon; 4.1% sodium cresol-formaldehyde sulphonated condensate; 8.0% urea; 0.7% polyalkylene glycol ether and polyoxyethylene alkylaryl ether; 0.5% silicic antifoaming agent; 0.2% xanthan gum; 0.5% of a 38.0% formaldehyde solution and 45.4% water.

17. A composition according to Claim 12, comprising, on a weight to volume basis: 30.0% pendimethalin; 5.0% of the water soluble ammonium salt of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid; 4.0% sodium lignosulphonate; 8.0% ethylene glycol; 0.5% silicic antifoaming; 0.75% silica; 0.1% Xanthan gum; 0.25% of a 38.0% formaldehyde solution; and 58.1% water.

18. A composition according to Claim 12, comprising, on a weight to volume basis: 20.0% pendimethalin; 30.0% chlortoluron; 3.6% sodium cresol-formaldehyde sulphonated condensate; 8.0% urea; 40 0.55% polyalkylene glycol ether and polyoxyethylene alkylaryl ether; 0.1% silicic antifoaming agent; 0.1% silica; 0.2% xanthan gum; 0.5% of a 38% formaldehyde solution; and 43.8% water.

19. A composition according to Claim 12, comprising, on a weight to volume basis: 26.0% pendimethalin; 12.5% of an isomeric mixture of methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluate and methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-p-toluate; 3.0% sodium cresol-formaldehyde condensate; 1.3% triethanolamine salt of polyarylethoxylate phosphate; 6.0% alkylphenolethoxylate; 5.0% ethylene glycol; 0.4% silicic antifoaming agent; 2.0% silica; 0.1% Xanthan gum; 0.25% of a 38% formaldehyde solution; and 49.62% water.

20. A method for preparing a stable aqueous suspension concentrate composition of pendimethalin alone or in combination with one or more secondary pesticides, said method comprising: emulsifying molten pendimethalin in hot water, temperature 50°C to 80°C, containing a surfactant and antifoaming agent, wherein said pendimethalin-surfactant-antifoaming agent mixture has a droplet size of about 2 microns to 10 microns; adding a suspending agent; and cooling the hot emulsion to ambient temperature, while agitating.

21. A method according to Claim 20, wherein said secondary pesticide(s) is water soluble or has a 55 melting point greater than 70°C.

22. A method according to Claim 21, additionally comprising: milling said emulsion following said cooling and agitation.

23. A method according to Claim 22, wherein said molten pendimethalin is emulsified in hot water containing surfactant, antifoaming agent and one or more pesticide(s) having a melting point greater than 70°C.

24. A method according to Claim 22, additionally comprising: adding said second pesticide(s) with melting point greater than 70°C and coformulants following said cooling of said emulsion to ambient temperature.

25. A method according to Claim 22, wherein said surfactant is an ethylene oxide/propylene oxide condensate; alkyl,aryl-, or aryl,aryl-ethoxylate and derivatives thereof; lignosulfonates; cresol-formaldehyde condensates and sulfonates; naphthalene-formaldehyde condensates and sulfonates; polycarboxylates and derivatives thereof; and mixture thereof.

26. A method according to Claim 25, wherein said suspending agent is a polysaccharide gum.

27. A method according to Claim 26, wherein said surfactant is a cresol formaldehyde condensate or sulfonate thereof; naphthalene-formaldehyde condensate or sulfonate thereof; a polycarboxylate or derivative thereof; a lignosulfonate; and mixtures thereof.

28. A method according to Claim 27, wherein said suspending agent is Xanthan gum.

29. A method according to Claim 28, wherein said composition is 30% to 40% pendimethalin; 3.0% to 5.0% sodium cresol-formaldehyde condensate or sodium cresol-formaldehyde sulphonated condensate surfactant; 5% to 10% ethylene glycol or urea antifreezing agent; 0.1% to 1.0% silicic antifoaming agent; 0.1% to 0.3% Xanthan gum suspending agent; 0.2% to 10% preservative of a 38% formaldehyde solution, methyl or propyl parahydroxybenzoate, 2-bromo-2-nitro-propane-1,3-diol, sodium benzoate, glutaraldehyde, O-phenylphenol, benzisothiazolinones, 5-chloro-2-methyl-4-isothiazolin-3-one, pentachlorophenol, 2-4-dichlorobenzylalcohol, or mixtures thereof; and 48.3% to 68.3% water.

30. A method according to Claim 28, wherein said composition is 40.0% pendimethalin; 5.0% sodium cresol-formaldehyde sulphonated condensate surfactant; 8.0% ethylene glycol; 0.5% silicic antifoaming agent; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution preservative; and 51.4% water.

31. A method according to Claim 28, wherein said composition is 27.3% pendimethalin; 18.2% atrazine; 3.4% sodium cresol-formaldehyde sulphonated condensate surfactant; 8.0% urea; 1.34% of polyalkylene glycol ether and poly oxyethylene alkylaryl ether surfactants; 0.9% silicic antifoaming agent; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution preservative; and 48.9% water.

32. A method according to Claim 28, wherein said composition is 23.6% pendimethalin; 23.6% isoproturon; 4.1% sodium cresol-formaldehyde sulphonated condensate surfactant; 8.0% urea; 0.7% polyalkylene glycol ether and polyoxyethylene alkylaryl ether surfactant; 0.5% silicic antifoaming agent; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution preservative; and 45.4% water.

33. A method according to Claim 28, wherein said composition is 30.0% pendimethalin; 5.0% of the water soluble ammonium salt of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid second pesticide; 4.0% sodium lignosulphonate; 8.0% ethylene glycol antifreezing agent; 0.5% silicic antifoaming agent; 0.75% silica antifoaming agent; 0.1% xanthan gum suspending agent; 0.25% of a 38.0% formaldehyde solution preservative; and 58.1% water.

34. A method according to Claim 28, wherein said composition is 20.0% pendimethalin; 30.0% chlortoluron second pesticide; 3.6% sodium cresol-formaldehyde sulphonated condensate surfactant; 8.0% urea; 0.55% polyalkylene glycol ether and polyoxyethylene alkylaryl ether surfactants; 0.1% silicic antifoaming agent; 0.1% silica; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution preservative; and 43.8% water.

35. A method according to Claim 28, wherein said composition is 23.6% pendimethalin; 23.6% isoproturon; 4.1% sodium cresol-formaldehyde sulphonated condensate surfactant; 8.0% urea; 0.7% blend of polyoxyethylene alkylaryl ether surfactant; 0.5% silicic antifoaming agent; 0.2% Xanthan gum; 0.5% of a 38% formaldehyde solution preservative; and 45.4% water.

36. A method for the preparation of a stable aqueous suspension concentrate composition of pendimethalin, said method comprising: dispersing molten pendimethalin in water, at ambient temperatures, containing surfactant and antifoaming agent; adding a suspending agent, 0% to 50.0%, on a weight to volume basis, of one or more secondary pesticide(s) and additional coformulants of dispersing agents, wetting agents, antifreezing agents, thickening agents, and preservatives; and milling to obtain a particle size of 20 microns.

37. A method according to Claim 36, wherein said secondary pesticide(s) is water soluble or has a melting point greater than 70°C.

38. A method according to Claim 37, wherein said surfactants, dispersing agents and wetting agents are ethylene oxide/propylene oxide condensates; alkyl,aryl-and aryl,aryl-ethoxylates; lignosulfonates; cresol-formaldehyde condensates and sulfonates; naphthalene-formaldehyde condensates and sulfonates; polycarboxylates and their derivatives; and mixtures thereof.

39. A method according to Claim 38, wherein said suspending agents are polysaccharide gums and cellulose derivatives.

40. A method according to Claim 39, wherein said polysaccharide gums are Xanthan gum, guar gum, gum arabic, and mixtures thereof.

41. A method according to Claim 39, wherein said antifreezing agents are ethylene glycol, propylene glycol, glycerine, urea, and mixtures thereof.

42. A method according to Claim 41, wherein said thickening agents are clays, precipitated silicas, polyvinyl alcohol, polyvinyl pyrrolidone, polyacrylamides, and mixtures thereof.

43. A method according to Claim 42, wherein said pesticide with a melting point greater than 70°C is Isoproturon, [N,N-dimethyl-N'-(4-(1-methylethyl)phenyl)urea]; Linuron, [N-3,4-dichlorophenyl-N'-methoxy-N'-methyl urea]; Metoxuron, [N'-(3-chloro-4-methoxyphenyl)-N,N-dimethylurea]; Chlortoluron, [N'-(3-chloro-4-methylphenyl)-N,N-dimethylurea]; Atrazine, [2-chloro-4-ethylamino-6-isopropylaminol,3-5-triazine]; Imidazolinone herbicides such as 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid and water soluble salts thereof or the isomeric mixture of methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluate and methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-toluate.

44. A method according to Claim 43, wherein said antifoaming agent is a silicic antifoaming agent.

45. A method according to Claim 44, wherein said preservatives is a 38% formaldehyde solution.

46. A composition according to Claim 11, comprising on a weight to volume basis: 23.6% pendimethalin, 23.6% isoproturon; 4.1% sodium cresol-formaldehyde sulphonated condensate; 0.71% polyalkylene glycol ether and polyoxyethylene alkylaryl ether; 8.0% urea; 0.2% xanthan gum; 0.5% silicic antifoaming agent; 0.1% methyl paraben; 0.05% propyl paraben; and 39.14% water.

47. A composition according to Claim 11, comprising on a weight to volume basis: 20.0% pendimethalin; 30.0% chlortoluron; 4.14% sodium cresol-formaldehyde sulphonated condensate; 1.64% polycarboxylate; 8.0% urea; 0.2% xanthan gum; 0.3% silicic antifoaming agent; 0.033% benzisothiazolinone and 35.68% water.

48. A method for preparing a stable aqueous suspension concentrate composition of pendimethalin alone or in combination with one or more secondary pesticides, said method comprising: emulsifying molten pendimethalin in hot water, temperature 50°C to 80°C, containing a surfactant, antifoaming agent and suspending agent while agitating; milling said resulting mixture whilst cooling, temperature 20°C to 25°C; aging the resulting cooled mixture, 0.5 hour to 48.0 hours; adding a secondary pesticide if desired; milling a second time; adding preservatives, surfactants, suspending agents, antifreezing agents.

49. A method according to Claim 48, wherein said surfactant is an ethylene oxide/propylene oxide condensate; alkyl,aryl-, or aryl,aryl-ethoxylate and derivatives thereof; lignosulfonates; cresol-formaldehyde condensates and sulfonates; naphthalene-formaldehyde condensates and sulfonates; polycarboxylates and derivatives thereof; and mixture thereof.

50. A method according to Claim 49, wherein said suspending agent is polysaccharide gum.

51. A method according to Claim 50, wherein said preservative is 0.2% to 10% of a 38% formaldehyde solution methyl or propyl parahydroxybenzoate, 2-bromo-2-nitro-propane-1,3-diol, sodium benzoate, glutaraldehyde, O-phenylphenol, benzisothiazolinones, 5-chloro-2-methyl-4-isothiazolin-3-one, pentachlorophenol, 2,4-dichlorobenzylalcohol, or mixtures thereof.



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X,P	EP-A-0 190 995 (CIBA-GEIGY) * Claims *	1-12	A 01 N 33/18 A 01 N 25/04

X	EP-A-0 163 598 (CIBA-GEIGY) * Claims; page 6, lines 4-6 *	1-19	

A,D	DE-A-3 302 648 (HOECHST) * Claims; page 7, line 21 - page 8, line 7 *	1-19	

A	FR-A-2 332 053 (BAYER)	20,36	

A	FR-A- 932 716 (MURPHY CHEMICAL)	20,36	

A,D	EP-A-0 033 291 (RHONE-POULENC) * Page 5, line 34 - page 6, line 34 *	48	TECHNICAL FIELDS SEARCHED (Int. Cl.4) A 01 N

The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 25-09-1987	Examiner DECORTE D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	